

WARNING. KEEP OUT OF REACH OF CHILDREN

ECOTOXIC

DONAGHYS

Myco-RF[®]

A systemic fungicide for the control of rust and Fusarium spp. in pasture and crown rust, stem rust and blind seed in ryegrass seed crops.

Active Ingredient: Contains 800 g/kg tebuconazole in the form of a wettable powder.



1kg

Registered pursuant to the ACVM Act 1997, No P007625
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996. Approval Code: HSR007810

DONAGHYS
PERFORMANCE FIRST

DONAGHY'S Myco-RF®

Read label before use.

PRODUCT INFORMATION: Myco-RF® has both contact and systematic action. It is absorbed by the leaves and stems and transported in an upwards and outwards movement in the plant, which provides protection into new growth. Full coverage is therefore necessary as plant parts below the area of contact will not be protected. It is an **offence to use this product on animals.**

COMPATIBILITY: Myco-RF® is compatible with most commonly used herbicides and insecticides. Myco-RF® should be used in conjunction with Donaghys Mycowet® Surfactant.

DIRECTIONS FOR USE

Recommended use rate: Apply 236g Myco-RF® per hectare

Mixing: Each 1kg pack is sufficient to treat 4.2 hectares of pasture. Open foil pack and remove inner pack which is soluble in water. Add unopened inner pack to partly filled spray tank with agitation running and then top up with water.

Rainfall: Do not apply if rain is likely within 1 hour of application. If 15mm or more of rain occurs within 3 hours of treatment, then pasture should be resprayed.

PASTURE TREATMENT FOR RUST AND FUSARIUM SPP

Boom Spraying: Apply 236g Myco-RF® plus 100ml Donaghys Mycowet® Surfactant in 100 litres of water per hectare. Increase Mycowet® to 150ml per hectare with water rates of 101-150 litres per hectare and to 200ml per hectare with water rates of 151-200 litres per hectare.

Aerial Spraying: Apply 236g Myco-RF® plus 100ml Donaghys Mycowet® Surfactant in 50 litres of water per hectare. Apply at first sign of rust development and Fusarium spp. A further application may be necessary later if climatic conditions favour re-infection.

Ensure complete coverage of all areas likely to be grazed. Spray should not be applied when air temperature exceeds 20°C or wind velocity exceeds 12kph.

CROWN RUST AND STEAM RUST

Rye Grass, Stem Rust: Apply 236g Myco-RF® plus 100ml Donaghys Mycowet® Surfactant in a minimum of 200 litres water/ha at the first appearance of the disease. A further application may be necessary 4 weeks later if disease pressure remains high or re-infection occurs.

Blind Seed Disease: Apply the above rates and make the first application at start of flowering and a second 10-14 days later.

NOTE: Do not apply Myco-RF® to the same pasture area or crop at less than 28 day intervals or more than two times per season.

REGULATORY STATEMENTS: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD

Grass Seed Crops: 30 days.

Pasture: Animals may be reintroduced into pasture treated with Myco-RF® 10 days after application. Animals for slaughter must be placed on a clean pasture for 6 days prior to slaughter. Pasture is considered clean from residues 45 days after treatment. When pasture cannot be guaranteed clean such as in aerial applications animals must not be sent for slaughter for 51 days after treatment of the pasture.

FACIAL ECZEMA PREVENTION: Myco-RF® does not control the fungus *Phythomyces* Chaptalia which is the causal organism for Facial Eczema. Myco-RF® is however compatible with Donaghys LIQUID Mycotak® (Carbendazim) and Donaghys Mycowet® Surfactant and

can be applied together where the control of rust (*Puccinia* spp) Fusarium spp. and *P. chaptalia* is sought. When applying together the same water and surfactant rates apply.

ENVIRONMENTAL WARNING: Very toxic to aquatic life. Avoid release to the environment. Collect spillage. Myco-RF® shall not be applied onto or into water.

PRECAUTIONS: Do not eat, drink or smoke while using, do not breathe dust, fumes, gas, mist or vapour spray. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Use only outdoors or in a well-ventilated area.

HEALTH WARNING: Harmful if swallowed, inhaled or comes in contact with skin. Causes mild skin irritation. May cause damage to organs through prolonged or repeated exposure.

FIRST AID

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTRE or doctor/physician if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water. Wash contaminated clothing before reuse. If skin irritation occurs: Get medical advice/attention.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Call a POISON CENTRE or doctor/physician if you feel unwell.

If medical advice is needed, have product container or label at hand. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or doctor immediately.

STORAGE: Store in original container, tightly closed in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. Storage must be in accordance with the current version of NZS8409 Management of Agrichemicals. This product is classified under HSDN 6.1 and 9.1A, and storage of certain quantities may require bunding and signage. See the SDS for further information. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product older than this.

DISPOSAL: When empty, triple rinse container and add residues into spray tank. Dispose of empty container at an approved landfill or other approved facility. Do not re-use container. Avoid contamination of any water supply with product or empty container.

NOTICE TO BUYER

Exclusion of liability: To the extent permitted by law, buyers and/or users of Myco-RF® accept that neither the manufacturer or Donaghys as distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of Myco-RF®.

Net Contents

1 kg

National Poisons Centre
0800 POISON (0800 764 766)



Acute oral toxicity Category 4. Acute dermal toxicity Category 4. Acute inhalation toxicity Category 4. Specific target organ toxicity - repeated exposure Category 2. Hazardous to the aquatic environment chronic Category 1. Hazardous to terrestrial vertebrates.

MANUFACTURED FOR AND DISTRIBUTED BY:
DONAGHY'S LTD, 16 SHEFFIELD CRESCENT
PO BOX 20 449, CHRISTCHURCH
FREEPHONE: 0800 942 006, WWW.DONAGHY'S.COM



DONAGHY'S
PERFORMANCE FIRST