

DANGER

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

TrivAL

Sheep HiMin



TRIPLE ACTIVE COMBINATION ORAL DRENCH FOR SHEEP

For the treatment and control of all major gastrointestinal parasites in sheep, including strains with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families. Also treats and controls lungworm, itch mite, tapeworm, larval stages of nasal bot, and aids in the control of adult liver fluke infection in sheep. Provides a supplemental source of copper, **selenium**, zinc, iodine and cobalt.

NET CONTENTS

20 Litres

ACTIVE INGREDIENTS

Abamectin 1 g/L; Levamisole Hydrochloride 40 g/L;
Albendazole 25 g/L; Copper 2.1 g/L;
Selenium 0.5 g/L; Zinc 0.6 g/L;
Cobalt 0.21 g/L and Iodine 0.76 g/L.

DONAGHYS
PERFORMANCE FIRST

DONAGHYS

TrivAL

Sheep HiMin

READ ENTIRE LABEL BEFORE USE

Harmful if inhaled. May cause allergic skin reactions. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs.

HANDLING PRECAUTIONS: Avoid breathing spray or mist. Do not eat, drink or smoke when handling this product. Contaminated clothing should not be allowed out of the workplace. Wear protective gloves. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid contact during pregnancy/ while nursing. Wash hands thoroughly after handling. Avoid release to the environment.

FIRST AID: If medical advice needed, have container or label at hand. IF SWALLOWED: Call a POISON CENTER (0800 764 766) or doctor/physician if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs get medical advice/attention. Wash contaminated clothing before reuse. IF exposed or concerned: Get medical advice/attention.

ENVIRONMENTAL PROTECTION: Very toxic to aquatic life. Hazardous to terrestrial vertebrates. Hazardous to terrestrial invertebrates.

Donaghys TrivAL Sheep HiMin contains albendazole, a member of the benzimidazole family, levamisole, a member of the imidazothiazole family and abamectin, a member of the macrocyclic lactone family. It is effective against gastrointestinal worms sensitive to one, two or all three broadspectrum drench families.

INDICATIONS:

SHEEP: For the effective treatment and control of mature and immature stages of susceptible strains of the following gastrointestinal parasites (* includes inhibited fourth stage larvae): *Teladorsagia circumcincta**, *Teladorsagia trifurcata**, *Haemonchus contortus**, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*, *Cooperia curticei**, *Cooperia oncophora**, *Cooperia punctata**, *Trichuris ovis*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Strongyloides papillosus*, *Nematodirus filicollis* and *Nematodirus spathiger*. Also effective against tapeworm (*Moniezia expansa*), lungworm (*Dictyocaulus filaria*), itch mite (*Psorobia ovis*), nodule worm (*Oesophagostomum columbianum*) and all larval stages of nasal bot (*Oestrus ovis*) and as an aid in the control of adult liver fluke (*Fasciola hepatica*) infestations in sheep.

Anthelmintic resistant strains: For the effective treatment and control of those gastrointestinal worms that have single and dual resistance to this combination of benzimidazole, levamisole/ morantel or avermectin/milbemycin anthelmintic action families. Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

DIRECTIONS FOR USE:


By law the user must take due care, obtaining expert advice when necessary to avoid unnecessary pain and distress when using this product other than directed on the label.

SHAKE WELL BEFORE USE. For oral administration only. Ensure formulation remains evenly suspended during use. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight, then dose according to the weight of the heaviest animal in the line. When there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Do not underdose. Correct drenching technique should be used. Check accuracy of drench guns regularly. Rinse gun with water immediately after use.

Accuracy of drench guns should be checked before and during drenching. Ensure that the dosing technique is accurate and that each sheep receives and swallows the full dose.

DOSE RATE: 1 mL per 5 kg bodyweight equivalent to albendazole 5 mg, levamisole hydrochloride 8 mg, abamectin 0.2 mg, selenium 0.1 mg per kg bodyweight.

DRUM TAP ENCLOSED.
Fit to internal thread after removing small drum cap.



Cut along dotted line

Bodyweight	Dose	Doses per 20L Pack
15 kg	3.0 mL	6666
16-20 kg	4.0 mL	5000
21-25 kg	5.0 mL	4000
26-30 kg	6.0 mL	3333
31-35 kg	7.0 mL	2857
36-40 kg	8.0 mL	2500
41-45 kg	9.0 mL	2222
46-50 kg	10.0 mL	2000
51-55 kg	11.0 mL	1818
56-60 kg	12.0 mL	1666
61-65 kg	13.0 mL	1538
66-70 kg	14.0 mL	1428
71-75 kg	15.0 mL	1333
Over 75 kg add 1 mL for every 5 kg bodyweight		

WARNINGS AND CONTRAINDICATIONS:

- Do not use on animals which are severely stressed or in ill health;
- Do not exceed stated dose or frequency without consulting veterinarian;
- Do not use in lambs under 15kg or under 4 weeks of age;
- Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate live weights carefully. Dehydrated animals may be more susceptible to toxicity;
- Do not treat sheep at intervals of less than 3 weeks, due to selenium content;
- Do not use at the same time as any other selenised fertilizer, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian;
- In sheep, liver levels of copper may be quite variable. Consequently there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species;
- Fatal interactions may occur between levamisole and organophosphate dips.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of the last treatment;
- Milk intended for human consumption must be discarded during treatment and for not less than 35 days following last treatment.

STORAGE: Store locked up below 25°C, out of direct sunlight.

SHELF LIFE: Product is suitable for use up to 18 months from the date of production (see batch number, expiry date) or within 12 months post-opening, whichever comes first.

DISPOSAL: Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at approved landfill facility. Do not use container for any other purpose.

Registered pursuant to the ACVM Act 1997: A011629.

See www.foodsafety.govt.nz for registration conditions.

EPA Approval number HSR100758.

See www.epa.govt.nz for approval conditions

REGISTERED TO

Donaghys Ltd, 16 Sheffield Crescent
Harewood, Christchurch, New Zealand

Freephone: 0800 942 006

www.donaghys.com

DONAGHYS

TrivAL

Sheep HiMin



UN3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE
(Contains Abamectin and Selenium)
Packaging Group III
HazChem 3Z

 **AGRECOVERY**
www.agrecovery.co.nz

Batch:

Expiry:

ZPLAB961 MAY25 PN-037722-1 VER3