

TrivAL[®]

Tape HiMin

TRIPLE ACTIVE COMBINATION PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control of mature stages of all major gastrointestinal parasites in sheep, including those with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families. Also treats and controls lungworm, itch mite, nasal bot, tapeworm, and as an aid in the control of adult liver fluke infestation in sheep. Also provides a supplemental source of **selenium**, copper, cobalt, zinc, and iodine.

NET CONTENTS
10 Litres

DANGER
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE
FOR ANIMAL TREATMENT ONLY

ACTIVE INGREDIENTS
Abamectin 1 g/L; Levamisole Hydrochloride 40 g/L; Albendazole 25 g/L; Praziquantel 19 g/L; **Selenium (as sodium selenate) 0.5 g/L**; Copper (as copper sulphate) 2.1 g/L; Zinc (as zinc sulphate) 0.6 g/L; and Iodine (as ethylenediamine dihydroiodide) 0.76 g/L.

DONAGHYS
PERFORMANCE FIRST

TrivAL[®]

Tape HiMin

TRIPLE ACTIVE COMBINATION PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control of mature stages of all major gastrointestinal parasites in sheep, including those with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families. Also treats and controls lungworm, itch mite, nasal bot, tapeworm, and as an aid in the control of adult liver fluke infestation in sheep. Also provides a supplemental source of **selenium**, copper, cobalt, zinc, and iodine.

NET CONTENTS
10 Litres

DANGER
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE
FOR ANIMAL TREATMENT ONLY

ACTIVE INGREDIENTS
Abamectin 1 g/L; Levamisole Hydrochloride 40 g/L; Albendazole 25 g/L; Praziquantel 19 g/L; **Selenium (as sodium selenate) 0.5 g/L**; Copper (as copper sulphate) 2.1 g/L; Zinc (as zinc sulphate) 0.6 g/L; and Iodine (as ethylenediamine dihydroiodide) 0.76 g/L.

DONAGHYS
PERFORMANCE FIRST

READ ENTIRE LABEL BEFORE USE

Donaghys TrivAl Tape HiMin contains albendazole, a member of the benzimidazole family, levamisole, a member of the imidazothiazole family and abamectin, a member of the macrocyclic lactone family. It is effective against gastrointestinal worms sensitive to one, two or all three broad spectrum drench families. It also contains praziquantel, a member of the isoquinoline family which is effective against tapeworm. Provides a supplemental source of selenium, copper, cobalt, zinc and iodine.

INDICATIONS Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

Sheep: For the effective treatment and control of mature and immature stages of the following gastrointestinal parasites (* includes inhibited fourth stage larvae):

*Teladorsagia circumcincta**, *Teladorsagia trifurcata**, *Haemonchus contortus**, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*, *Cooperia curticei**, *Cooperia oncophora**, *Cooperia punctata**, *Trichuris ovis*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Strongyloides papillosus*, *Nematodirus filicollis* and *Nematodirus spathiger*.

Also effective against all stages of tapeworm (*Moniezia expansa*), lungworm (*Dictyocaulus filaria*), itch mite (*Psorobia ovis*), nodule worm (*Oesophagostomum columbianum*) and all larval stages of nasal bot (*Oestrus ovis*) and as an aid in the control of adult liver fluke infestations in sheep.

Anthelmintic resistant strains: For the effective treatment and control of those gastrointestinal worms that have single and dual resistance to this combination of benzimidazole, levamisole/morantel or avermectin/milbemycin anthelmintic action families.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label

SHAKE WELL BEFORE USE. For oral administration only. Ensure formulation remains evenly suspended during use. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight, then dose according to the weight of the heaviest animal in the line. When there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Do not underdose. Correct drenching technique should be used. Check accuracy of drench guns regularly. Rinse gun with water immediately after use. Accuracy of drench guns should be checked before and during drenching. Ensure that the dosing technique is accurate and that each sheep receives and swallows the full dose.

DOSE RATE

1 mL per 5 kg bodyweight equivalent to albendazole 5 mg, levamisole hydrochloride 8 mg, abamectin 0.2 mg, praziquantel 3.8 mg, selenium 0.1 mg, copper 0.42 mg, cobalt 0.04 mg, zinc 0.12 mg and iodine 0.15 mg per kg bodyweight.

DOSE RATE

Bodyweight:	Dose	Doses per 10L pack
20 kg	4 mL	2500
21-25 kg	5 mL	2000
26-30 kg	6 mL	1666
31-35 kg	7 mL	1428
36-40 kg	8 mL	1250
41-45 kg	9 mL	1111
46-50 kg	10 mL	1000
51-55 kg	11 mL	909
56-60 kg	12 mL	833
61-65 kg	13 mL	769
66-70 kg	14 mL	714
71-75 kg	15 mL	666
Over 75 kg add 1 mL for every 5 kg bodyweight		

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment;

Milk intended for human consumption must be discarded during treatment and for not less than 35 days following last treatment.

STORAGE

Store locked up below 25°C, out of direct sunlight.

Registered to: Donaghys Ltd,
16 Sheffield Crescent, PO Box 20 449, Christchurch
Freephone: 0800 942 006 www.donaghys.com

EPA Approval number HSR100758.

See www.epa.govt.nz for approval conditions
Registered pursuant to the ACVM Act 1997: A011205.
See www.foodsafety.govt.nz for registration conditions.



Causes serious eye irritation. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause an allergic skin reaction. Suspected of causing cancer. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs through prolonged or repeated exposure.

HANDLING PRECAUTIONS: Do not handle until all safety precautions have been read and understood. Do not breathe mist/vapours/spray. In case of inadequate ventilation wear respiratory protection. Do not eat, drink, or smoke when using this product. Take off contaminated clothing and wash before reuse. Contaminated clothing should not be allowed of the workplace. Wear protective gloves/protective clothing/eye protection. Use personal protective equipment as required. Avoid contact during pregnancy and while nursing. Wash hands thoroughly after handling.

FIRST AID: If inhaled: Remove person to fresh air and keep comfortable for breathing. If experiencing respiratory symptoms: Call the National Poisons Centre 0800 POISON (0800 764 766) or doctor/physician. If exposed or if you feel unwell or concerned: Get medical advice/attention. If medical advice is needed, have container or label at hand. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. If on skin: Wash with plenty of soap and water. If skin irritation or rash occurs get medical advice/attention.

ENVIRONMENTAL PROTECTION: Very toxic to aquatic life with long lasting effects. Hazardous to terrestrial vertebrates. Hazardous to terrestrial invertebrates. Avoid release to the environment. Collect spillage.

DISPOSAL: Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at approved landfill facility. Do not use container for any other purpose.

EMERGENCY MANAGEMENT: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

WARNINGS AND CONTRAINDICATIONS

- Do not use on animals that are severely stressed or in ill health
- Do not exceed stated dose or frequency without consulting veterinarian
- Do not use in lambs under 20 kg or under 6 weeks of age
- Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate live weights carefully. Dehydrated animals may be more susceptible to toxicity
- Do not treat sheep at intervals of less than 3 weeks, due to selenium content
- Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian
- In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species
- Fatal interactions may occur between levamisole and organophosphate dips.

DRUM TAP ENCLOSED.
Fit to internal thread after removing small drum cap.



Cut along dotted line.



ENVIRONMENTALLY HAZARDOUS SUBSTANCE
(Contains Abamectin, Selenium)

ZPLA9860 JULY25 FN 038120-1 VER 4