

WARNING

KEEP OUT OF REACH
OF CHILDREN
READ LABEL
BEFORE USE
FOR ANIMAL
TREATMENT ONLY

DONAGHYS

MoxiPor

POUR ON



LONG LASTING ENDECTOCIDE FOR CATTLE AND DEER

For the treatment and control of internal and external parasites of cattle (including lactating dairy cattle) and for the treatment and control of lungworm and roundworms of deer.

Improves milk production in lactating dairy cows and increases weight gains in growing beef and dairy cattle exposed to parasite challenge.

NET CONTENTS
20 Litres

ACTIVE CONSTITUENT
Moxidectin 5g/L
Contains Aromatic Hydrocarbons 150g/L

DONAGHYS
PERFORMANCE FIRST

DONAGHYS

MoxiPor

POUR ON

LONG LASTING ENDECTOCIDE
FOR CATTLE AND DEER

DONAGHYS

MoxiPor

POUR ON

LONG LASTING ENDECTOCIDE
FOR CATTLE AND DEER

READ LABEL CAREFULLY BEFORE USING Obtain special instructions before use. Do not handle it until all safety precautions have been read and understood.

HANDLING PRECAUTIONS Combustible liquid. Keep away from heat, hot surfaces. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. Do not breathe mist, vapours or spray. Avoid skin contact. Avoid contact during pregnancy and while nursing. Handle with care. Wear protective gloves and use personal protective equipment, as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse. Store locked up.

FIRST AID For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If exposed or concerned, get medical advice/attention.

ENVIRONMENTAL PROTECTION Very toxic to the aquatic life with long-lasting effects. Avoid contamination of any water supply with product or empty container. Hazardous to soil organisms. Hazardous to terrestrial invertebrates. Avoid release to the environment.

EMERGENCY RESPONSE In case of fire, use water fog, dry chemical powder of CO2 to extinguish. In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further material. Dispose of waste safely.

DISPOSAL Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse the empty container, crush or puncture and bury in a suitable landfill, or if appropriate, recycle. Do not use container for any other purpose.

MOXIPOR POUR ON contains moxidectin, a member of the milbemycin/ avermectin family. It is effective against sensitive strains of the following internal parasites:

CATTLE **Gastrointestinal Roundworms:** Adult and immature forms unless otherwise stated; *Haemonchus contortus* (barber's pole worm), *Ostertagia ostertagi* (*O. lyrata*) (small brown stomach worm including inhibited larvae), *Trichostrongylus axei* (stomach hair worm), *Trichostrongylus colubriformis* (black scour worm), *Trichostrongylus longispicularis*, *Cooperia oncophora* (small intestinal worm), *Cooperia punctata*, *Oesophagostomum radiatum* (nodule worm), *Bunostomum phlebotomum* (hookworm).

Lungworm: Adult and immature *Dictyocaulus viviparus*. **Lice:** *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola bovis*. **Mange mites:** *Chorioptes bovis*.

DEER **Lungworm:** *Dictyocaulus ekerti* (viviparus). **Gastrointestinal roundworms:** *Haemonchus contortus*, *Ostertagia*-like species*, *Trichostrongylus* spp, *Oesophagostomum venulosum*, *Chabertia ovina* **Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera* *assymetrica*, *Spiculoptera* *spiculoptera* and *Skrjabinagia* *kolchida*.

PERSISTENT ACTIVITY **MOXIPOR POUR ON**, following a single application at the recommended dose rate, prevents reinfection of:

CATTLE with *Dictyocaulus viviparus*, *Oesophagostomum radiatum* and *Bunostomum phlebotomum* for 42 days, *Ostertagia ostertagi* (*O. lyrata*) for 35 days and *Haemonchus* spp and *Trichostrongylus axei* for 28 days, and **DEER** with *Dictyocaulus ekerti* (viviparus), *Ostertagia*-like spp*, *Oesophagostomum venulosum* and *Chabertia ovina* for 42 days. **Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera* *assymetrica*, *Spiculoptera* *spiculoptera* and *Skrjabinagia* *kolchida*.

Resistance may develop to any anthelmintic. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.

DRUM TAP ENCLOSED.
Fit to internal thread after removing small drum cap.



Cut along dotted line

DOSAGE AND ADMINISTRATION:
Dose Rate: 1 mL/10 kg to give 0.5 mg moxidectin per kg liveweight.

Liveweight (kg)	Dose Volume (mL)	Doses per 20L Pack	Liveweight (kg)	Dose Volume (mL)	Doses per 20L Pack
50	5	4000	250	25	800
60	6	3333	300	30	666
70	7	2857	350	35	571
80	8	2500	400	40	500
90	9	2222	450	45	444
100	10	2000	500	50	400
110	11	1818	550	55	363
120	12	1666	600	60	333
130	13	1538	650	65	307
140	14	1428	700	70	285
150	15	1333	750	75	266
200	20	1000			

Registered pursuant to the ACVM Act 1997, No. A11390
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act, No HSR100759
See <http://www.epa.govt.nz> for controls.

DIRECTIONS FOR USE:
By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. **MOXIPOR POUR ON** should be applied along the top of the back from the shoulders to the base of the tail. Check accuracy of the pour-on applicator regularly. A representative sample of animals should be weighed before treatment either with scales or a weighband. Choose dose according to the weight of the heaviest animals. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight to avoid excessive overdosing.

Rainfast: The efficacy of **MOXIPOR POUR ON** is not adversely affected if applied when the hide is wet, or if rain occurs shortly after the application. However treatment of animals under these circumstances is not recommended practice.

Non-irritant: **MOXIPOR POUR ON** does not cause irritation to the skin or damage to the hide when used as directed.

Ease of Use: **MOXIPOR POUR ON** comes ready to use in applicator bottles, back packs or drums.

Concurrent Use: **MOXIPOR POUR ON** is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines. **It should not be applied at the same time as any other pour-on product.**

Safety to Target Animals: **MOXIPOR POUR ON** has a wide margin of safety when used at the recommended dose rate.

WITHHOLDING PERIODS:
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT
Cattle - Nil
Deer - Nil
Bobby Calves - Nil
Milk - Nil

STORAGE: Store locked up below 25°C, in a well ventilated place and protect from light.

DONAGHYS

Registered to:
Horizon Agresources (NZ) Limited

Marketed and Distributed by:
Donaghys Ltd
16 Sheffield Crescent,
Harewood, Christchurch, New Zealand
Freephone: 0800 942 006
www.donaghys.com



UN3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.
(Contains Moxidectin 0.5%)
Packaging Group III



Batch:
Expiry:

ZPLAB37 SEP25 PN-039010-1 VER2