

DONAGHYS

ProAbamec

POUR-ON FOR CATTLE

For the treatment and control of internal and external parasites of cattle.

ACTIVE INGREDIENTS
Abamectin 10 mg/ml
Contains >941 g/L
Diethylene glycol monobutyl ether

NET CONTENTS
5 Litre

DONAGHYS
PERFORMANCE FIRST

WARNING
FLAMMABLE LIQUID
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

DONAGHYS

ProAbamec

READ ENTIRE LABEL BEFORE USE

WARNING: Dangerous to the environment.

HANDLING PRECAUTIONS Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Keep away from heat/sparks/open flames/hot surfaces. No smoking. Do not breathe fume/gas/mist/vapours/spray. Use personal protective equipment as required. Wear protective gloves/ protective clothing/ eye protection/face protection. Avoid contact during pregnancy/while nursing. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area.

HAZARD STATEMENTS Harmful if swallowed, inhaled or absorbed through the skin. May cause mild skin and eye irritation. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. Repeated exposure may cause skin allergy. Abamectin may possibly affect development and/or reproduction and may have effects on/or via lactation. Diethylene glycol monobutyl ether may possibly affect kidney if swallowed and liver if inhaled.

ENVIRONMENTAL PROTECTION Very toxic to aquatic life. Hazardous to soil organisms. Hazardous to terrestrial vertebrates. Hazardous to terrestrial invertebrates. Avoid release to the environment. Collect spillage.

FIRST AID If swallowed: Rinse mouth. Do NOT induce vomiting. If on skin: Wash with plenty of soap and water. Wash contaminated clothing before reuse. If inhaled: Remove to fresh air and keep at rest in a position comfortable for breathing. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. If exposed or concerned: Get medical advice/ attention. If medical advice is needed, have product container or label at hand. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

EMERGENCY RESPONSE In case of spillage wear appropriate protective clothing. Prevent material from entering waterways and absorb spills with inert material and place in waste container. Wash area and absorb with further inert material.

INDICATIONS Donaghys Proabamec Pour-On for Cattle, when used at the recommended dose level of 500 micrograms Abamectin per kg liveweight, effectively controls the following parasites of cattle:

GASTRO-INTESTINAL ROUNDWORMS Adult and immature *Ostertagia spp* (including inhibited immatures), *Haemonchus spp*, *Trichostrongylus spp*, *Cooperia spp*, *Bunostomum spp*, *Oesophagostomum spp*, and adults of *Chabertia spp*, *Nematodirus spp*, *Trichuris spp*, and *Strongyloides spp*.

LUNGWORMS Adult and immature *Dictyocaulus viviparus*.

BITING AND SUCKING LICE *Bovicola bovis*, *Linognathus vituli*.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Donaghys ProAbamec Pour-On for Cattle contains 10 mg/ml Abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family. Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Check accuracy of pour-on applicator regularly. A representative sample of animal should be weighed before treatment. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc). Do not under-dose. Where there is a large variation in size within the group, draft

into two or more lines based on bodyweight, to avoid excessive overdosing. Ready to use with standard pour-on equipment. Apply along the backline from shoulders to tail.

Donaghys ProAbamec Pour-On is rainfast, even if rain falls 2 hours or more after treatment.

DOSAGE It is strongly recommended not to use this product in cattle under four months of age and/or 100 kg live weight. The dose rate for cattle is 500µg Abamectin per kg bodyweight, which is provided by 1 mL Donaghys ProAbamec Pour-On for Cattle per 20 kg bodyweight, or according to the table:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

MILK Milk intended for sale for human consumptions must be discarded during treatment and for **35 days** following the last treatment.

MEAT Cattle producing meat and offal for human consumption must not be sold for slaughter during or within **35 days** of the last treatment.

STORAGE Store locked up, below 25°C. Protect from sunlight and keep it well closed.

DISPOSAL Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.

Registered pursuant to the ACVM Act 1997 No.A010875.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, approval code HSR100759.
See www.epa.govt.nz for controls.

REGISTERED TO AND DISTRIBUTED BY
Donaghys Ltd, 16 Sheffield Crescent, Harewood, Christchurch
Freephone: 0800 942 006
www.donaghys.com

UN No.3082, Packaging Group III, HazChem 3Z
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Contains Abamectin)

Batch:
Expiry:

DOSE RATE		
Live-weight (kg)	Dose (mL)	5L Pack Treats
101-150 kg	7.5 mL	666
151-200 kg	10 mL	500
201-250 kg	12.5 mL	400
251-300 kg	15 mL	333
301-350 kg	17.5 mL	285
351-400 kg	20 mL	250
401-450 kg	22.5 mL	222
451-500 kg	25 mL	200
501-550 kg	27.5 mL	181
551-600 kg	30 mL	166
601-650 kg	32.5 mL	153
Heavier animals (e.g. mature bulls) should receive an additional 1 mL for each 20 kg over 650 kg.		

